



**BeneFix**<sup>®</sup>

Coagulation Factor IX (Recombinant)

Room Temperature Storage

\*BeneFix was approved February 11, 1997.



Not actual patients.

# EXPERIENCE MATTERS

BeneFix: The only recombinant factor IX supporting individuals with hemophilia for more than 20 years.\*

## BeneFix has dosing flexibility that may help fit your lifestyle

- Indicated for bleed control and prevention in individuals with hemophilia B
- The convenience of the BeneFix Rapid Reconstitution (R2) Kit with a range of vial sizes
- Safety profile demonstrated in clinical trials



## What Is BeneFix?

BeneFix, Coagulation Factor IX (Recombinant), is an injectable medicine that is used to help control and prevent bleeding in people with hemophilia B. Hemophilia B is also called congenital factor IX deficiency or Christmas disease.

BeneFix is **NOT** used to treat hemophilia A.

## Important Safety Information

- BeneFix is contraindicated in patients who have manifested life-threatening, immediate hypersensitivity reactions, including anaphylaxis, to the product or its components, including hamster protein.
- Call your health care provider right away if your bleeding is not controlled after using BeneFix.
- Allergic reactions may occur with BeneFix. Call your health care provider or get emergency treatment right away if you have any of the following symptoms: wheezing, difficulty breathing, chest tightness, your lips and gums turning blue, fast heartbeat, facial swelling, faintness, rash or hives.
- Your body can make antibodies, called "inhibitors," which may stop BeneFix from working properly.

Reach out to your local Patient Affairs Liaison to learn more about Pfizer Hemophilia's products, resources, and support

Linda Pollhammer  
302-258-5985 [linda.pollhammer@pfizer.com](mailto:linda.pollhammer@pfizer.com)

- If you have risk factors for developing blood clots, such as a venous catheter through which BeneFix is given by continuous infusion, BeneFix may increase the risk of abnormal blood clots. The safety and efficacy of BeneFix administration by continuous infusion have not been established.
- Some common side effects of BeneFix are nausea, injection site reaction, injection site pain, headache, dizziness and rash.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see the Brief Summary for BeneFix on the next page.



R<sub>x</sub> only

### Brief Summary

See package insert for full Prescribing Information. This product's label may have been updated. For further product information and current package insert, please visit [www.Pfizer.com](http://www.Pfizer.com) or call our medical communications department toll-free at 1-800-438-1985.

Please read this Patient Information carefully before using BeneFix and each time you get a refill. There may be new information. This brief summary does not take the place of talking with your doctor about your medical problems or your treatment.

### What is BeneFix?

BeneFix is an injectable medicine that is used to help control and prevent bleeding in people with hemophilia B. Hemophilia B is also called congenital factor IX deficiency or Christmas disease.

BeneFix is **NOT** used to treat hemophilia A.

### What should I tell my doctor before using BeneFix?

Tell your doctor and pharmacist about all of the medicines you take, including all prescription and non-prescription medicines, such as over-the-counter medicines, supplements, or herbal remedies.

Tell your doctor about all of your medical conditions, including if you:

- are pregnant or planning to become pregnant. It is not known if BeneFix may harm your unborn baby.
- are breastfeeding. It is not known if BeneFix passes into the milk and if it can harm your baby.

### How should I infuse BeneFix?

The initial administrations of BeneFix should be administered under proper medical supervision, where proper medical care for severe allergic reactions could be provided.

### See the step-by-step instructions for infusing in the complete patient labeling.

You should always follow the specific instructions given by your doctor. If you are unsure of the procedures, please call your doctor or pharmacist before using.

### Call your doctor right away if bleeding is not controlled after using BeneFix.

Your doctor will prescribe the dose that you should take. Your doctor may need to test your blood from time to time. BeneFix should not be administered by continuous infusion.

### What if I take too much BeneFix?

Call your doctor if you take too much BeneFix.

### What are the possible side effects of BeneFix?

Allergic reactions may occur with BeneFix. Call your doctor or get emergency treatment right away if you have any of the following symptoms:

wheezing	fast heartbeat
difficulty breathing	swelling of the face
chest tightness	faintness
turning blue (look at lips and gums)	rash
	hives

Your body can also make antibodies, called "inhibitors," against BeneFix, which may stop BeneFix from working properly.

Some common side effects of BeneFix are nausea, injection site reaction, injection site pain, headache, dizziness and rash.

BeneFix may increase the risk of thromboembolism (abnormal blood clots) in your body if you have risk factors for developing blood clots, including an indwelling venous catheter through which BeneFix is given by continuous infusion. There have been reports of severe blood clotting events, including life-threatening blood clots in critically ill neonates, while receiving continuous-infusion BeneFix through a central venous catheter. The safety and efficacy of BeneFix administration by continuous infusion have not been established.

These are not all the possible side effects of BeneFix.

Tell your doctor about any side effect that bothers you or that does not go away.

### How should I store BeneFix?

DO NOT FREEZE BeneFix. The BeneFix kit can be stored at room temperature (below 86°F) or under refrigeration. Throw away any unused BeneFix and diluent after the expiration date indicated on the label.

Freezing should be avoided to prevent damage to the pre-filled diluent syringe.

BeneFix does not contain a preservative. After reconstituting BeneFix, you can store it at room temperature for up to 3 hours. If you have not used it in 3 hours, throw it away.

Do not use BeneFix if the reconstituted solution is not clear and colorless.

### What else should I know about BeneFix?

Medicines are sometimes prescribed for purposes other than those listed here. Do not use BeneFix for a condition for which it was not prescribed. Do not share BeneFix with other people, even if they have the same symptoms that you have.

If you would like more information, talk to your doctor. You can ask your doctor for information about BeneFix that was written for healthcare professionals.

This brief summary is based on BeneFix® [Coagulation Factor IX (Recombinant)] Prescribing Information LAB-0464-10.0, revised June 2017.